



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

I. Solution by H. M. ARMSTRONG, Cooch's Bridge, Delaware.

Let the sides of the given triangle be denoted by the vectors 2α , 2β , and $2(\alpha + \beta)$. Let i denote a unit vector perpendicular to the plane of the triangle. Then

$$\rho_1 = \alpha + \frac{1}{\sqrt{3}}i\alpha \dots\dots\dots (1), \quad \rho_2 = 2\alpha + \beta + \frac{1}{\sqrt{3}}i\beta \dots\dots\dots (2), \quad \rho_3 = \alpha + \beta - i(\alpha + \beta) \dots\dots\dots (3)$$

are the vectors drawn to the vertices of the three isosceles triangles.

Hence the vectors of the sides of the new triangle are obviously $\rho_2 - \rho_1$, $\rho_3 - \rho_2$, $\rho_1 - \rho_3$. By means of (1), (2), and (3), we find that $(\rho_2 - \rho_1)^2 = (\rho_3 - \rho_2)^2 = (\rho_1 - \rho_3)^2$.

Hence $T(\rho_2 - \rho_1) = T(\rho_3 - \rho_2) = T(\rho_1 - \rho_3)$. Therefore the triangle is equilateral.

II. Solution by G. W. GREENWOOD, M. A. (Oxon). Lebanon, Ill.

Denote the vertices by A , B , C , and the remaining vertices of the triangles adjacent to BC , CA , AB , by A' , B' , C' , respectively. I will assume that the triangles are all exterior to the given triangle. Then

$$\begin{aligned} A'C &= a/\sqrt{3}, \quad B'C = b/\sqrt{3}, \quad \angle A'CB' = 60^\circ + C. \\ A'B'^2 &= \frac{1}{3}[a^2 + b^2 - 2ab\cos(60^\circ + C)] \\ &= \frac{1}{3}[a^2 + b^2 - ab(\cos C - \sqrt{3}\sin C)] \\ &= \frac{1}{6}[a^2 + b^2 + c^2 + 2\sqrt{3} ab\sin C] \\ &= \frac{1}{6}[a^2 + b^2 + c^2 + \sqrt{3} abc/r], \end{aligned}$$

where r is the radius of the circumscribed circle of the triangle ABC . The result shows that $A'B' = B'C' = C'A'$. The same result will follow if A' and A are on the same side of BC , etc.

Also solved by M. E. Graber, J. Scheffer, and G. B. M. Zerr. G. I. Hopkins refers to solution of Problem 96, Vol. 5, No. 4.

245. Proposed by J. H. M. MACLAGAN-WEDDERBURN, M. A., Chicago, Ill.

Given two lines of fixed length, $AB = a$, $BC = b$, perpendicular to each other. A line CP is drawn making $\angle BPC = \theta$. AD is the perpendicular to AB meeting CP in D . Find by Euclidean construction the angle θ such that $AD^2 \cos^2 \theta + b^2 \sin^2 \theta$ is a minimum.

Solution by the PROPOSER.

Draw any line CP . From B drop the perpendicular BL upon CP ; draw AM parallel to PC meeting BL in M . Join CM . Then $CM^2 = CL^2 + LM^2 = b^2 \sin^2 \theta + AD^2 \cos^2 \theta$. The locus of M is a circle described on AB as a diameter with O , the middle point of AB , as center. In order that CM^2 , and therefore CM shall be a minimum, the line CM must pass through O .

The construction is therefore the following: Bisect AB in O ; produce BOA to P so that $OP = OC$. Then $BPC =$ required angle θ . [Newton's *Principia*, Vol. 2, Section 7, Proposition XXXIV, Scholium.]